

LOW-MASS X-RAY BINARIES, MILLISECOND RADIO PULSARS, AND THE COSMIC STAR FORMATION RATE

Nicholas E. White & Pranab Ghosh^{1,2}

Laboratory for High Energy astrophysics

Code 662, NASA Goddard Space Flight Center, Greenbelt, MD 20771

¹ Senior NAS/NRC Resident Research Associate

² On leave from Tata Institute of Fundamental Research, Bombay 400 005, India

Received 8th June 1998; accepted 2nd July 1998

To appear in the Astrophysical Journal Letters

ABSTRACT

We report on the implications of the peak in the cosmic star-formation rate (SFR) at redshift $z \approx 1.5$ for the resulting population of low-mass X-ray binaries (LMXB) and for that of their descendants, the millisecond radio pulsars (MRP). Since the evolutionary timescales of LMXBs, their progenitors, and their descendants are thought to be significant fractions of the time-interval between the SFR peak and the present epoch, there is a lag in the turn-on of the LMXB population, with the peak activity occurring at $z \sim 0.5 - 1$. The peak in the MRP population is delayed further, occurring at $z \lesssim 0.5$. We show that the discrepancy between the birthrate of LMXBs and MRPs, found under the assumption of a steady-state SFR, can be resolved for the population as a whole when the effects of a time-variable SFR are included. A discrepancy may persist for LMXBs with short orbital periods, although a detailed population synthesis will be required to confirm this. Further, since the integrated X-ray luminosity distribution of normal galaxies is dominated by X-ray binaries, it should show strong luminosity evolution with redshift. In addition to an enhancement near the peak ($z \approx 1.5$) of the SFR due to the prompt turn-on of the relatively short-lived massive X-ray binaries and young supernova remnants, we predict a second enhancement by a factor ~ 10 at a redshift between ~ 0.5 and ~ 1 due to the delayed turn-on of the LMXB population. Deep X-ray observations of galaxies out to $z \approx 1$ by AXAF will be able to observe this enhancement, and, by determining its shape as a function of redshift, will provide an important new method for constraining evolutionary models of X-ray binaries.

Subject headings: Binaries: close – Galaxies: evolution – pulsars: general – stars: evolution – stars: formation – X-rays: galaxies – X-rays: stars

1. Introduction

It has recently been shown that the cosmic star formation rate (SFR) increases with redshift, reaching a peak ~ 10 times higher than the current rate in the redshift interval $\sim 1 - 2$ (Lilly *et al.* 1996; Madau *et al.* 1997; Madau, Pozzetti & Dickinson 1998, henceforth MPD; Madau, della Valle & Panagia 1998, henceforth MVP). In this *Letter*, we report interesting implications of this for the evolution of LMXBs, for that of their descendants, the millisecond radio pulsars (MRP), and also for the well-known LMXB-MRP “birthrate problem” (see Bhattacharya 1995 and references therein). Similar considerations of the implications of the cosmic SFR for the evolution of the cosmic supernova rates have recently been undertaken (MVP). In essence, the LMXB-MRP birthrate problem stems from the observation that, for estimated numbers $N_{LMXB} \sim 100$ and $N_{MRP} \sim 10^4$ in our galaxy, and for expected lifetimes $\tau_{LMXB} \sim 10^8 - 10^9$ yr and $\tau_{MRP} \sim 10^9 - 10^{10}$ yr, the steady-state birthrate of MRPs, $\frac{N_{MRP}}{\tau_{MRP}}$, exceeds that of LMXBs, $\frac{N_{LMXB}}{\tau_{LMXB}}$, by $\gtrsim 10$ (Kulkarni and Narayan 1988, henceforth KN; Côté & Pylyser 1991, henceforth CP; Lorimer 1995, henceforth L95). Many suggested solutions to the problem include (a) accretion-induced collapse of a white dwarf to a neutron star, (b) much shorter values of τ_{LMXB} for the short-period LMXBs (*i.e.*, those with orbital periods $\lesssim 3$ days, say, following the definition of KN), due, *e.g.*, to X-ray irradiation of the low-mass companion (Tavani 1991, henceforth T91), and (c) the possibility that pulsars can be born with low magnetic fields and millisecond periods.

We show here that steady-state arguments do not generally apply to evolving LMXB and MRP populations with a time-dependent global SFR peaking at $z \approx 1.5$, since this peak propagates through the LMXB and MRP populations at smaller redshifts, and causes an enhanced MRP population at the present epoch. Indeed, except in special circumstances outlined in §2, there is no basis for expecting an equality between the rates $\frac{N_{MRP}}{\tau_{MRP}}$ and $\frac{N_{LMXB}}{\tau_{LMXB}}$. We present evolutionary calculations which show that the expected

number ratio, $N_r \equiv \frac{n_{MRP}}{n_{LMXB}}$, and rate ratio, $R_r \equiv \frac{N_{MRP}}{\tau_{MRP}} / \frac{N_{LMXB}}{\tau_{LMXB}}$, at $z = 0$ are in agreement with the currently observed values for the whole population. However, there may still be a discrepancy for the short-period systems. We consider other observational tests of our model and indicate how observations of galaxies in the redshift range $0.5 - 1.0$ by AXAF can constrain models for the evolution of X-ray binaries.

2. Evolution of LMXBs & MRPs With Variable Star Formation Rate

The standard evolutionary scenario for the majority¹ of LMXBs and MRPs begins with a primordial binary containing a massive OB and a low-mass star (see, *e.g.*, Webbink, Rappaport & Savonije 1983, henceforth WRS; Kalogera & Webbink, 1996, 1998). The massive star rapidly evolves to the point of supernova (SN), resulting in the formation of a post-SN binary (PSNB) consisting of a neutron star with a low-mass companion, which turns into a LMXB when the latter attains Roche lobe contact, either due to nuclear evolution or due to orbital decay by gravitational radiation and magnetic braking. This, in turn, produces a recycled MRP at the end of mass transfer. We demonstrate here the basic effects of a time-variable SFR on the above scenario. In this introductory work, we confine ourselves to a simple description, in which the evolution of the number density of each species (n_{PSNB} for PSNBs, n_{LMXB} for LMXBs, and n_{MRP} for MRPs) is described by a timescale which is a given number: τ_{PSNB} for the evolution of PSNBs into LMXBs, τ_{LMXB} for that of LMXBs into MRPs, and τ_{MRP} for that of the MRPs. In reality, τ_{PSNB} and τ_{LMXB} depend on the binary period, τ_{PSNB} on other evolutionary parameters (see below)

¹We do not consider the LMXBs and MRPs found in globular clusters, where tidal capture of neutron stars in close encounters with stars causes an excess of LMXBs relative to the overall Galactic population.

as well, and τ_{MRP} on evolutionary parameters which seem poorly understood at present. To demonstrate the effects of these timescales, we run the evolutionary scheme over the ranges of their values suggested in the literature: detailed population-synthesis studies are deferred to the future.

The evolutions of populations of PSNBs, LMXBs and MRPs in response to a time-dependent star-formation rate $SFR(t)$ are given by:

$$\frac{\partial n_{PSNB}(t)}{\partial t} = \alpha SFR(t) - \frac{n_{PSNB}(t)}{\tau_{PSNB}}, \quad (1)$$

$$\frac{\partial n_{LMXB}(t)}{\partial t} = \frac{n_{PSNB}(t)}{\tau_{PSNB}} - \frac{n_{LMXB}(t)}{\tau_{LMXB}}, \quad (2)$$

$$\frac{\partial n_{MRP}(t)}{\partial t} = \frac{n_{LMXB}(t)}{\tau_{LMXB}} - \frac{n_{MRP}(t)}{\tau_{MRP}}. \quad (3)$$

In equation (1), $SFR(t)$ is that given by MPD and MVP, with the SFR evolving on a timescale $\tau_{SFR} \approx 6.4 \times 10^8$ yr: for all calculations reported here, we have used the analytic approximation (accurate to within 5%) to the SFR given by MVP, which is shown in Figure 1. Further, α is a coefficient which determines the rate of formation of PSNBs per unit star-formation rate. Assuming that the time required by massive newborn stars to evolve to the point of supernova is small compared to all other evolutionary timescales in the problem, *i.e.*, τ_{SFR} , τ_{PSNB} , τ_{LMXB} , and τ_{MRP} (an excellent approximation in view of the value of τ_{SFR} given above and the values of τ_{PSNB} , τ_{LMXB} , and τ_{MRP} given below), α is given approximately by $\alpha = \frac{1}{2} f_{binary} f_{prim} f_{SN}$. Here, f_{binary} is the fraction of all stars in binaries, f_{prim} is that fraction of primordial binaries which has the correct range of stellar masses and orbital periods for evolving into PSNBs capable of producing LMXBs (KW98), and f_{SN} is that fraction of the latter binaries which survives the supernova. The actual value of α , which sets the overall scale for the sizes of the populations n_{PSNB} , n_{LMXB} and n_{MRP} relative to that of the SFR, is irrelevant for this study, since we are only interested in the relative sizes of n_{LMXB} and n_{MRP} here.

Figure 1 shows the evolution of PSNBs, LMXBs and MRPs described by equations (1)-(3): we have displayed our results in terms of the redshift z , which is related to the cosmic time t by $t_9 = 13(z + 1)^{-3/2}$, where t_9 is t in units of 10^9 yr, and a value of $H_0 = 50$ km s $^{-1}$ Mpc $^{-1}$ has been used (MPD). Our choices of representative values of τ_{PSNB} , τ_{LMXB} , and τ_{MRP} come from the following considerations. The distribution of τ_{PSNB} with orbital period is rather broad, has a peak in the range $1 - 2 \times 10^9$ yr, and is somewhat dependent on supernova kick velocity and common-envelope evolution efficiency (Kalogera 1998, private communication). The expected range of τ_{LMXB} values has been discussed extensively in the literature on the birthrate problem (KN; CP; T91; L95). The standard WRS mass-transfer time, $\tau_{LMXB} \approx 1.1 \times 10^9 / P_i(d)$ yr has been widely used, where $P_i(d)$ is the initial orbital period of the LMXB in days, and, to ameliorate the problem for short-period LMXBs, it has been suggested that the effects of X-ray irradiation of the low-mass companion may reduce τ_{LMXB} to $\sim 10^7$ yr in these systems (T91; L95). For τ_{MRP} , we used values based on the compilation by Camilo, Thorsett & Kulkarni (1994), which suggests τ_{MRP} in the range $3 \times 10^9 - 3 \times 10^{10}$ yr.

3. Discussion

The results shown in Figures 1(a)-(c) clearly demonstrate that the $\sim$ Gyr timescales involved in the evolution of PSNBs, LMXBs and MRPs lead to substantial time-lags in the peaks of their populations behind the peak in the SFR. The LMXB peak is delayed by several Gyr relative to the SFR peak, and appears in the redshift range $z \sim 0.5 - 1$. The MRP peak is delayed even further, appearing at redshifts $\lesssim 0.5$ (including the current epoch). Thus, although the previous work of KN and others assumed steady state conditions while comparing the birthrates of MRP and LMXB, we now see that this assumption is not correct in general in a universe with time-dependent cosmic SFR. There

are, however, two limiting cases where this assumption will still apply. The first occurs when an asymptotic state is reached for all populations, which happens at times much longer than *all* evolutionary timescales in the problem, as illustrated by the case shown in Figure 1(d). However, it must be clear that such a situation cannot occur within the current age of the universe for any realistic choice of τ_{PSNB} , τ_{LMXB} , and τ_{MRP} , which is why we had to assume unrealistically short timescales for Figure 1(d). This possibility is thus of little relevance to the present universe, unless our current understanding of LMXB and MRP evolution is completely wrong. The second situation obtains when, for sufficiently large values of τ_{MRP} , the present epoch ($z=0$) happens to be at or near the maximum of the MRP evolution, where $\frac{\partial n_{MRP}(t)}{\partial t} = 0$ (see eq.[3]), so that $\frac{n_{MRP}}{\tau_{MRP}} \approx \frac{n_{LMXB}}{\tau_{LMXB}}$ at this epoch. As illustrated by the case shown in Figure 1(a), such a situation is quite possible for realistic values of evolutionary timescales. We have demonstrated in Figures 1(b) and (c) that it is also possible have situations in which $\frac{n_{MRP}}{\tau_{MRP}}$ is considerably larger than $\frac{n_{LMXB}}{\tau_{LMXB}}$ in the present epoch for plausible values of evolutionary timescales.

In relating the observational situation to the basic theoretical expectations for the number ratio, N_r , and rate ratio, R_r of evolving LMXB and MRP populations, we first emphasize that $\frac{n_{MRP}}{\tau_{MRP}}$ and $\frac{n_{LMXB}}{\tau_{LMXB}}$ are really the instantaneous rates of *decay* of the MRP and LMXB populations, respectively (see eqs. [2]-[3]), and not their “birthrates”, as they have been often called in previous discussions. Only under the assumption of a steady state can we equate them to the respective birthrates, and to each other. For evolving populations, $R_r \approx 1$ is expected only under the circumstances described in the last paragraph. Thus, there is no basis for expecting $\frac{n_{MRP}}{\tau_{MRP}} = \frac{n_{LMXB}}{\tau_{LMXB}}$ in general, and a deviation from equality does not, by itself, imply a serious problem. Indeed, since the observable quantities are really n_{MRP} and n_{LMXB} , the actual test of agreement is as follows: given a plausible choice of τ_{PSNB} , τ_{LMXB} and τ_{MRP} , does the calculated number ratio N_r at $z = 0$ at the present epoch agree with observation, and, furthermore, does the calculated rate ratio R_r at the

same epoch agree with that obtained from the observed N_r with this particular choice of timescales?

With the discovery of many more MRPs since the original KN work, the observational situation has changed somewhat. KN estimated N_r to be $\sim 10^2$, and R_r to be ~ 10 for the whole population and ~ 100 for short-period systems. The most recent estimate by L95 (using ~ 5 times as many MRPs as KN) suggests $N_r \approx 400$, $R_r \approx 1$ for the whole population, and $R_r \approx 8$ for short-period systems. From the case in Figure 1(a), it is clear that values of N_r and R_r typically discussed for the whole population are naturally obtained in the above picture with canonical timescales for the whole population. If short-period systems with longer τ_{LMXB} (KN; CP) are considered alone, Figure 1(b) shows that $R_r > 1$ also occurs naturally, but with typical R_r values ≈ 3 . It is not clear how significant the discrepancy for short-period LMXBs is until a more detailed population synthesis has been undertaken. If further work confirms the discrepancy, we may conclude that either (a) a one-to-one evolution from PSNB to LMXB to MRP does not always occur: in certain parts of the parameter space, PSNBs do not evolve into the LMXB phase, but ultimately produce MRPs (KW98), (b) some potential LMXBs are rapidly destroyed, possibly by evaporation of the secondary (CP; T91), or, (c) there is a serious undersampling of the LMXB population because the majority are not X-ray active (L95).

4. An Observational Test

We have demonstrated the inadequacy of steady-state arguments in discussions of the LMXB-MRP birthrate problem for evolving populations with a time-variable SFR that peaks at a redshift ≈ 1.5 . We find that an evolutionary scheme can easily account for the observed MRP/LMXB number ratios, *i.e.*, $R_r \approx 1$ for the overall population, and larger R_r values for short-period LMXB systems. A closely related point is the relative

behavior of high-mass X-ray binary (HMXB) and LMXB populations in a universe with a time-dependent cosmic SFR. HMXBs and LMXBs originating from stars formed in the same epoch have very different evolutionary times, since, although the initial evolution of both involves the evolution of a massive star to supernova and the formation of a neutron star, LMXBs turn on as X-ray sources much later than HMXBs, only after the low-mass companion comes into Roche lobe contact, predominantly due to orbital decay by gravitational radiation and magnetic braking (KW98 and references therein). The relevant time lag is essentially the timescale τ_{PSNB} introduced in §2. Since the post-supernova evolution into HMXBs takes a negligibly short time on this scale, the global HMXB population will peak roughly where the number of stars (= integral of the SFR) does. Thus, the global LMXB population will peak in redshift well after the HMXB peak. The combined X-ray binary activity of the two populations is expected to have a broad peak, or possibly a double peak, in z , depending on the lag and the relative population sizes.

The dominant source of X-ray emission from normal spiral galaxies (*i.e.*, those without an active nucleus) appears to be the integrated emission from their X-ray binaries (see Fabbiano 1995 and references herein), based on observations of nearby galaxies such as M31 (where individual sources can be resolved) and comparison with the distribution in our galaxy. These integrated X-ray luminosities are in the range $L_x \sim 10^{39}$ – 10^{41} erg s $^{-1}$ and scale linearly with the blue band luminosities of the galaxies. In our galaxy, LMXB dominate the total X-ray output, and this is also the same for M31, where the brightest sources are clustered around the bulge. For other relatively nearby galaxies, the average X-ray temperatures are in the range 3 – 6 keV, also consistent with a population of LMXB (Kim, Fabbiano & Trinchieri 1992). HMXB systems are also a significant component in some galaxies. They dominate the 1–10 keV output of the irregular LMC and SMC galaxies. The X-ray outputs of starburst galaxies in the 1–10 keV band seem to be dominated by those of their HMXB populations and/or young supernova remnants (SNR; della Cecca,

Griffiths & Heckman 1997).

Our work demonstrates that the peak in the SFR at $z \approx 1.5$ will cause the integrated X-ray luminosity of galaxies in the redshift range $0.5 - 1.0$ to be at least an order of magnitude higher than it is today. If the current understanding of LMXB evolution is correct, then a twin-peak signature of the dual LMXB and HMXB-SNR population is expected. This is caused by the delayed turn-on of the LMXB population relative to the short-lived and instantaneous turn-on of the HMXBs and SNRs associated with the peak in the SFR (Figure 1). This second LMXB peak is in the redshift range $0.5 - 1.0$ and is caused by the delay of the secondary in the PSNB to come into contact with its Roche lobe. The details of this signature, *e.g.*, peak separation, can then be used to confirm the general picture as to the origin of LMXBs, and to constrain models for their evolution. The expected flux levels ($\sim 10^{-15}$ – 10^{-16} erg cm $^{-2}$ s $^{-1}$) for this redshift range will be within the capabilities of AXAF, provided sufficient observing time ($\sim 10^5$ – 10^6 s) is dedicated to a suitable field. These future observations will provide an important new window to understand the evolution of X-ray binaries and the resulting millisecond radio pulsar population.

It is a pleasure to thank V. Kalogera for communicating results of evolutionary calculations in advance of publication, and P. Madau for supplying the MVP approximation.

REFERENCES

- Bhattacharya, D. 1995, in X-ray Binaries, ed. W. H. G. Lewin, J. van Paradijs and E. P. J. van den Heuvel, 233
- Camilo, F., Thorsett, S.E. & Kulkarni, S.R. 1994, ApJ, 421, L15
- Coté, J., & Pylyser, E. H. P. 1989, A&A, 218, 131 (CP)
- della Cecca, R., Griffiths, R., & Heckman, T.M. 1997, ApJ, 485, 581
- Fabbiano, G. 1995, in X-ray Binaries, ed. W. H. G. Lewin, J. van Paradijs and E. P. J. van den Heuvel, 390
- Kalogera, V. & Webbink, R.F. 1996 ApJ, 458, 301
- Kalogera, V. & Webbink, R.F. 1998 ApJ, 493, 351 (KW98)
- Kim, D.W., Fabbiano, G., & Trinchieri, G. 1992 ApJ, 393, 134
- Kulkarni, S. & Narayan, R. 1988 ApJ, 335, 755 (KN)
- Lilly, S.J., Le Fèvre, O., Hammer, F., & Crampton, D. 1996, ApJ, 460, L1
- Lorimer, D. R. 1995, MNRAS, 274, 300 (L95)
- Madau, P., Ferguson, H. C., Dickinson, M. E., Giavalisco, M., Steidel, C. C., & Fruchter, A., 1996, MNRAS, 283, 1388
- Madau, P., Pozzetti, L., & Dickinson, M. 1998, ApJ, 498, 106 (MPD)
- Madau, P., della Valle, M., & Panagia, N. 1998, MNRAS, submitted (MVP)
- Tavani, M. 1991, ApJ, 366, L27 (T91)
- Webbink, R. F., Rappaport, S., & Savonije, G. J. 1983, ApJ, 270, 678 (WRS)




Fig. 1.— Evolution of PSNB, LMXB, and MRP populations in response to a time-variable cosmic star-formation rate (SFR). Shown are logarithms of the number densities of PSNBs (dotted line), LMXBs (dash-dotted line), and MRPs (solid line) against the redshift. The SFR of Madau *et al.* is also shown (dashed line) for reference. Each panel displays the results of an evolutionary calculation with input timescales τ_{PSNB} , τ_{LMXB} , and τ_{MRP} written at the top of the panel, and output values of the number ratio, $N_r \equiv \frac{n_{MRP}}{n_{LMXB}}$, and rate ratio, $R_r \equiv \frac{N_{MRP}}{\tau_{MRP}} / \frac{N_{LMXB}}{\tau_{LMXB}}$, at $z = 0$ written at the bottom. Case (a) represents a typical result for the whole population of LMXBs and MRPs, with $R_r \approx 1$. Case (b) represents a typical result for short-period systems (see text), with $R_r > 1$. Case (c) shows the results of postulating an unusually short LMXB lifetime, due, *e.g.*, to X-ray irradiation of the secondary in close binaries (see text). Finally, case (d) illustrates the approach to an asymptotic state for all populations, as described in the text, obtained by choosing unrealistically short values for the timescales τ_{PSNB} , τ_{LMXB} , and τ_{MRP} .